

AUTHORS Du,X., Poltorak,A., Wei,Y. and Beutler,B.
 TITLE Three novel mammalian toll-like receptors: gene structure,
 expression, and evolution
 JOURNAL Eur. Cytokine Netw. 11 (3), 362-371 (2000)
 PUBMED 11022119
 REFERENCE 7 (residues 1 to 1049)
 AUTHORS Rock,F.L., Hardiman,G., Timans,J.C., Kastelein,R.A. and Bazan,J.F.
 TITLE A family of human receptors structurally related to Drosophila Toll
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 95 (2), 588-593 (1998)
 PUBMED 9435236
 COMMENT REVIEWED REFSEQ: This record has been curated by NCBI staff. The
 reference sequence was derived from [AF245702.1](#) and [AF240467.1](#).

Summary: The protein encoded by this gene is a member of the
 Toll-like receptor (TLR) family which plays a fundamental role in
 pathogen recognition and activation of innate immunity. TLRs are
 highly conserved from Drosophila to humans and share structural and
 functional similarities. They recognize pathogen-associated
 molecular patterns (PAMPs) that are expressed on infectious agents,
 and mediate the production of cytokines necessary for the
 development of effective immunity. The various TLRs exhibit
 different patterns of expression. This gene is predominantly
 expressed in lung, placenta, and spleen, and lies in close
 proximity to another family member, TLR8, on chromosome X.

FEATURES Location/Qualifiers
 source 1..1049
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="X"
 /map="Xp22.3"
 Protein 1..1049
 /product="toll-like receptor 7"
 variation 11
 /replace="Q"
 /replace="L"
 /db_xref="dbSNP:179008"
 Region <75..>423
 /region_name="Membrane glycoprotein LIG-1 [Signal
 transduction mechanisms]"
 /note="KOG4194"
 /db_xref="CDD:21971"
 Region 256..768
 /region_name="Extracellular matrix protein slit, contains
 leucine-rich and EGF-like repeats [Extracellular
 structures, Signal transduction mechanisms]"
 /note="KOG4237"
 /db_xref="CDD:22013"
 Region 319..>833
 /region_name="Membrane glycoprotein LIG-1 [Signal
 transduction mechanisms]"
 /note="KOG4194"
 /db_xref="CDD:21971"
 variation 448
 /replace="A"
 /replace="V"
 /db_xref="dbSNP:5743781"
 Region 498..>654
 /region name="Ras suppressor protein (contains

Region

CDS

ORIGIN

```
1 mvfpmwtlkr qililfniil isklldgarwf pkltpcdvtl dvpknhvivd ctdkhlteip
61 ggipntntnl tltinhipdi spsfhrldh lveidfrnc vpiplgsknn mcikrlqikp
121 rsfsgltylk slyldgnqll eipqglppsl qllsleanni fsirkenlte lanieilylg
181 qncyyrnpcy vsysiekdaf lnltklkvls lkdnntavp tvlpstltel ylynnmiaki
241 qeddfnnlnq lqildlsgnc prcynapfpc apcknnsplq ipvnafalt elkvrlrlhsn
301 slqhvpprwf kninklqeld lsgnflakei gdakflhflp sliqldlsfn felqvyrasm
361 nlsqafsslk slkilrirgy vfkelksfnl splhnlqnle vldlgtfnik ianlsmfkqf
421 krlkvidlsv nkispsgdss evgfcnsart svesyepqvl eqlhyfrydk yarscrfknk
481 easfmsvnes cykygqtdl sknsiffvks sdfqhlslfk clnlsnlnis qtlngsefqp
541 laelryldfs nnrldllhst afeelhklef ldissnshyf qsegitmln ftknlkvlgk
601 lmmndndiss stsrtnses lrtlefrgnh ldvlwregdn rylqlfknll kleeldiskn
661 slsflpsgvf dgmpnlnknl slaknglksf swkklqclkn letldlshnq lttvperlsn
721 csrslnklil knngirsltk yflqdafqlr yldlssnkiq miqktsfpen vlennkmlll
781 hhnrlfctcd avwfvwvnh tevtipylat dvtevgpgah kgqsvisl dl ytceldltnl
841 ilfslsisvs lflmvmmtas hlyfwdvwyi yhfckakikg yqrlispdce ydafivdytk
901 dpavtewvla elvakledpr ekhfnlclee rdwlpqgpvl enlsqsiqls kktvfvmtdk
961 yaktenfkia fylshqrlmd ekvdviilif lekpfqkskf lqlrkrlcgs svlewptnpg
1021 aphyfwqcln nalatdnhva ysqvfketv
```

Sequence



Disclaimer | Write to the Help Desk
NCBI | NLM | NIH